

JEE ADVANCED-2017

PHYSICS

A. General Instructions :



1. This sealed booklet is your Question Paper. Do not break the seal till you are instructed to do so.
2. The question paper CODE is printed on the left hand top corner of this sheet and the right hand top corner of the back cover of this booklet
3. Use the Optical Response Sheet (ORS) provided separately for answering the questions.
4. The paper CODE is printed on its left part as well as the right part of the ORS. Ensure that both these codes are identical and same as that on the question paper booklet. If not, contact the invigilator.
5. Blank spaces are provided within this booklet for rough work.
6. Write your name and roll number in the space provided on the back cover of this booklet
7. After breaking the seal of the booklet at 9:00 am, verify that the booklet contains 36 pages and that all the 54 questions along with the options are legible. If not, contact the invigilator for replacement of the booklet
8. You are allowed to take away the Question Paper at the end of the examination.

Optical Response Sheet

9. The ORS (top sheet) will be provided with an attached Candidate's Sheet (bottom sheet). The Candidate's Sheet is a carbon - less copy of the ORS.
10. Darken the appropriate bubbles on the ORS by applying sufficient pressure. This will leave an impression at the corresponding place on the Candidate's Sheet
11. The ORS will be collected by the invigilator at the end of the examination.
12. You will be allowed to take away the Candidate's Sheet at the end of the examination.

13. Do not tamper with or mutilate the ORS. Do not use the ORS for rough work.
14. Write your name, roll number and code of the examination center, and sign with pen in the space provided for this purpose on the ORS. Do not write any of these details anywhere else on the ORS. Darken the appropriate bubble under each digit of your roll number.

Darken the Bubbles on the ORS

15. Use a Black Ball Point Pen to darken the bubbles on the ORS.
16. Darken the bubble  completely.
17. The correct way of darkening a bubble is as: 
18. The ORS is machine - gradable. Ensure that the bubbles are darkened in the correct way.
19. Darken the bubbles only if you are sure of the answer. There is no way to erase or "un-darken" a darkened bubble.

SECTION -1 : (Maximum Marks : 28)

- This section contains **SEVEN** questions.
- Each question has **FOUR** options (A), (B), (C) and (D), **ONE OR MORE THAN ONE** of these four option(s) is(are) correct.
- For each question, darken the bubble(s) corresponding to all the correct option(s) in the ORS.
- For each question, marks will be awarded in one of the following categories :

Full Marks +4 If only the bubble(s) corresponding to all the correct option(s) is(are) darkened.


Partial Marks +1 For darkening a bubble corresponding to each correct option, provided NO incorrect option is darkened.

Zero Marks 0 If none of the bubbles is darkened.


Negative Marks -2 In all other cases.

- For example, if (A), (C) and (D) are all the correct options for a question, darkening all these three will get +4 marks; darkening only (A) and (D) will get +2 marks and darkening (A) and (B) will get -2 marks, as a wrong option is also darkened.


1. A block M hangs vertically at the bottom end of a uniform rope of constant mass per unit length. The top end of the rope is attached to a fixed rigid support at O . A transverse wave pulse (Pulse 1) of wavelength λ_0 is produced at point O on the rope. The pulse takes time T_{OA} to reach point A . If the wave pulse of wavelength λ_0 is produced at point A (Pulse 2) without disturbing the position of M it takes time T_{AO} to reach point O . Which of the following options is/are correct ?



- (A) The velocities of the two pulses (Pulse 1 and Pulse 2) are the same at the midpoint of rope
- (B) The velocity of any pulse along the rope is independent of its frequency and wavelength
- (C) The wavelength of Pulse 1 becomes longer when it reaches point A
- (D) The time $T_{AO} = T_{OA}$
2. A circular insulated copper wire loop is twisted to form two loops of area A and $2A$ as shown in the figure. At the point of crossing the wires remain electrically insulated from each other. The entire loop lies in the plane (of the paper). A uniform magnetic field \vec{B} points into the plane of the paper. At $t = 0$, the loop starts rotation about the common diameter as axis with a constant angular velocity ω in the magnetic field. Which of the following options is/are correct?



- (A) The net emf induced due to both the loops is proportional to $\cos \omega t$
- (B) The rate of change of the flux is maximum when the plane of the loops is perpendicular to plane of the paper
- (C) The amplitude of the maximum net emf induced due to both the loops is equal to the amplitude of maximum emf induced in the smaller loop alone
- (D) The emf induced in the loop is proportional to the sum of the area of the two loops
3. A block of mass M has a circular cut with a frictionless surface as shown. The block rests on the horizontal frictionless surface of a fixed table. Initially the right edge of the block is at $x=0$, in a coordinate system fixed to the table. A point mass m is released from rest at the topmost point of the path as shown and it slides down. When the mass loses contact with the block, its position is x and the velocity is v . At the instant, which of the following options is/are correct?



(A) The velocity of the point mass m is : $v = \sqrt{\frac{2gR}{1 + \frac{m}{M}}}$

(B) The x component of displacement of the center of mass of the block M is :
 $-\frac{mR}{M+m}$

(C) The position of the point mass is $x = -\sqrt{2} \frac{mR}{M+m}$

(D) The velocity of the block M is: $V = -\frac{m}{M} \sqrt{2gR}$

4. A flat plate is moving normal to its plane through a gas under the action of a constant force F . The gas is kept at very low pressure. The speed of the plate v is much less than the average speed u of the gas molecules. Which of the following options is/are true?


(A) The pressure difference between the leading and trailing faces of the plate is proportional to uv .

(B) At a later time the external force F balances the resistive force

(C) The resistive force experienced by the plate is proportional to v

(D) The plate will continue to move with constant non-zero acceleration, at all times

5. In the circuit shown, $L = 1\mu H$, $C = 1\mu F$ and $R = 1k\Omega$. They are connected in series with an a.c. source $V = V_0 \sin \omega t$ as shown. Which of the following options is/are correct?



- (A) The current will be in phase with the voltage if $\omega = 10^4 \text{ rad.s}^{-1}$
- (B) At $\omega \gg 10^6 \text{ rad.s}^{-1}$, the circuit behaves like a capacitor
- (C) The frequency at which the current will be in phase with the voltage is independent of R
- (D) At $\omega \sim 0$ the current flowing through the circuit becomes nearly zero
6. For an isosceles prism of angle A and refractive index μ , it is found that the angle of minimum deviation $\delta_m = A$. Which of the following options is/are correct?
- (A) At minimum deviation, the incident angle i_1 and the refracting angle r_1 at the first refracting surface are related by $r_1 = \left(\frac{i_1}{2}\right)$
- (B) For this prism, the refractive index μ and the angle of prism A are related as $A = \frac{1}{2} \cos^{-1} \left(\frac{\mu}{2} \right)$
- (C) For the angle of incidence $i_1 = A$, the ray inside the prism is parallel to the base of the prism
- (D) For this prism, the emergent ray at the second surface will be tangential to the surface when the angle of incidence at the first surface is $i = \sin^{-1} \left[\sin A \sqrt{4 \cos^2 \frac{A}{2} - 1} - \cos A \right]$
7. A human body has surface area of approximately 1 m^2 . The normal body temperature is 10 K above the surrounding room temperature T_0 . Take the room temperature to be $T_0 = 300 \text{ K}$. For $T_0 = 300 \text{ K}$, the value of $\sigma T_0^4 = 460 \text{ Wm}^{-2}$ (where σ is the Stefan-Boltzmann constant). Which of the following options is/are correct ?

- (A) If the surrounding temperature reduces by a small amount $\Delta T_0 \ll T_0$, then to maintain the same body temperature the same (living) human being needs to radiate $\Delta W = 4\sigma T_0^3 \Delta T_0$ more energy per unit time.
- (B) Reducing the exposed surface area of the body (e.g. by curling up) allows humans to maintain the same body temperature while reducing the energy lost by radiation
- (C) If the body temperature rises significantly then the peak in the spectrum of electromagnetic radiation emitted by the body would shift to longer wavelengths
- (D) The amount of energy radiated by the body in 1 second is close to 60 joules

SECTION - 2 : (Maximum Marks : 15)

This section contains **FIVE** questions.

The answer to each question is a **SINGLE DIGIT INTEGER** ranging from 0 to 9, both inclusive. For each question, darken the bubble corresponding to the correct integer in the ORS.


For each question, marks will be awarded in one of the following categories :

Full Marks : +3 If only the bubble corresponding to the correct answer is darkened.

Zero Marks : 0 In all other cases.

8. A drop of liquid of radius $R = 10^{-2} \text{ m}$ having surface tension $S = \frac{0.1}{4\pi} \text{ Nm}^{-1}$ divides itself into K identical drops. In this process the total change in the surface energy $\Delta U = 10^{-3} \text{ J}$. If $K = 10^\alpha$ then the value of α is:

9. A monochromatic light is travelling in a medium of refractive index $n = 1.6$. It enters a stack of glass layers from the bottom side at an angle $\theta = 30^\circ$. The interfaces of the glass layers are parallel to each other. The refractive indices of different glass layers are monotonically decreasing as $n_m = n - m\Delta n$, where n_m is the refractive index of the m^{th} slab and $\Delta n = 0.1$ (see the figure). The ray is refracted out parallel to the interface between the $(m-1)^{\text{th}}$ and m^{th} slabs from the right side of the stack. What is the value m ?



10. ^{131}I is an isotope of Iodine that β decays to an isotope of Xenon with a half-life of 8 days..A small amount of a serum labelled with ^{131}I is injected into the blood of a person. The activity of the amount of ^{131}I injected was 2.4×10^5 Becquerel (Bq). It is known that the injected serum will get distributed uniformly in the blood stream in less than half an hour. After 11.5 hours, 2.5 ml of blood is drawn from the person's body, and gives an activity of 115 Bq. The total volume of blood in the person's body, in liters is approximately (you may use $e^x \approx 1 + x$ for $|x| \ll 1$ and $\ln 2 \approx 0.7$)
11. An electron in a hydrogen atom undergoes a transition from an orbit with quantum number n_i to another with quantum number n_f , V_i and V_f are respectively the initial and final potential energies of the electron. If $\frac{V_i}{V_f} = 6.25$, then the smallest possible n_f is:

12. A stationary source emits sound of frequency $f_0 = 492 \text{ Hz}$. The sound is reflected by a large car approaching the source with a speed of 2 ms^{-1} . The reflected signal is received by the source and superposed with the original. What will be the beat frequency of the resulting signal in Hz? (Given that the speed of sound in air is 330 ms^{-1} and the car reflects the sound at the frequency it has received).

SECTION - 3 : (Maximum Marks : 18)

- This section contains SIX questions of matching type,
- This section contains TWO tables (each having 3 columns and 4 rows).
- Based on each table, there are THREE questions
- Each question has FOUR options (A), (B), (C) and (D). ONLY ONE of these four options is correct,
- For each question, darken the bubble corresponding to the correct option in the ORS,
- For each question, marks will be awarded in one of the following categories :

Full Marks : +3 If only the bubble corresponding to the correct option is darkened.

Zero Marks : 0 If none of the bubbles is darkened.

Negative Marks : -1 In all other cases.

Answer Q.13. Q14 and Q.15 by appropriately matching the information given in the three columns of the following table.

A charged particle (electron or proton) is introduced at the origin ($x = 0, y = 0, z = 0$) with a given initial velocity \vec{v} . A uniform electric field \vec{E} and a uniform magnetic field \vec{B} exist everywhere. The velocity \vec{v} , electric field \vec{E} and magnetic field \vec{B} are given in column 1, 2 and 3, respectively. The quantities E_0, B_0 are positive in magnitude.

Column-1

Column-2

Column-3

(I) Electron with $\vec{v} = 2 \frac{E_0}{B_0} \hat{x}$

(i) $\vec{E} = E_0 \hat{z}$

(P) $\vec{B} = -B_0 \hat{x}$

(II) Electron with $\vec{v} = 2 \frac{E_0}{B_0} \hat{y}$

(ii) $\vec{E} = -E_0 \hat{y}$

(Q) $\vec{B} = B_0 \hat{x}$

(III) Electron with $\vec{v} = 0$

(iii) $\vec{E} = -E_0 \hat{x}$

(R) $\vec{B} = B_0 \hat{y}$

(IV) Electron with $\vec{v} = 2 \frac{E_0}{B_0} \hat{x}$

(iv) $\vec{E} = E_0 \hat{x}$

(S) $\vec{B} = B_0 \hat{z}$

13. In which case will the particle move in a straight line with constant velocity ?

(A) (IV) (i) (S)

(B) (III) (ii) (R)

(C) (II) (iii) (S)

(D) (III) (iii) (P)

14. In which case will the particle describe a helical path with axis along the positive z direction

(A) (IV) (i) (S)

(B) (II) (ii) (R)

(C)(III) (iii) (P)

(D) (IV) (ii) (R)

15. In which case would the particle move in a straight line along the negative direction of y -axis (i.e, move along $-\hat{y}$)

(A) (III) (ii) (R)

(B) (IV) (ii) (S)

(C) (III) (ii) (P)

(D) (II) (iii) (Q)

Answer Q.16. Q.17 and Q.18 by appropriately matching the information given in the three columns of the following table.

An ideal gas is undergoing a cyclic thermodynamic process in different ways as shown in the corresponding $P-V$ diagrams in column 3 of the table. Consider only the path from state 1 to state 2. W denotes the corresponding work done on the system. The equations and plots in the table have standard notations as used in thermodynamic process. Here γ is the ratio of heat capacities at constant pressure and constant volume. The number of moles in the gas is n .

Column-1

(I) $W_{1 \rightarrow 2} = -\frac{1}{\gamma-1}(P_2V_2 - P_1V_1)$



(II) $W_{1 \rightarrow 2} = -PV_2 + PV_1$

Column-2

(i) Isothermal


(ii) Isochoric

Column-3



(III) $W_{1 \rightarrow 2} = 0$


(iii) Isobaric



(R)

(IV) $W_{1 \rightarrow 2} = -nRT \ln \left(\frac{V_2}{V_1} \right)$

(iv) Adiabatic



(S)

16. Which of the following options is the only correct representation of a process in which $\Delta U = \Delta Q - P\Delta V$?

(A) (II) (iii) (P)

(B) (N) (iii) (S)

(C) (III) (Hi) (P)

(D) (II) (iv) (R)

17. Which one of the following options is the correct combination ?

(A) (II) (iv) (P)

(B) (IV) (ii) (S)

(C) (II) (iv) (R)

(D) (III) (ii) (S)

18. Which one of the following options correctly represents a thermodynamics process that is used as a correction in the determination of the speed of sound in an ideal gas ?

(A) (III) (iv) (R)

(B) (I) (H) (Q)

(C) (IV) (ii) (R)

(D) (I) (iv) (Q)

